

AD-A147 961

PHASE-DISSIPATIVE MECHANISMS FOR LASER-INDUCED SURFACE
DESORPTION/DISSOCI. (U) ROCHESTERUNIV NY DEPT OF
CHEMISTRY X Y HUANG ET AL. OCT 84

1/1


UNCLASSIFIED

UROCHESTER/DC/84/TR-56 N00014-80-C-0472

F/G 7/4

NL

END



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

12

AD-A147 961

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0472

Task No. NR 056-749

TECHNICAL REPORT No. 56

Phase-Dissipative Mechanisms for Laser-Induced Surface
Desorption/Dissociation Processes

by

Xi-Yi Huang, Thomas F. George, Jian-Min Yuan and L. M. Narducci

Prepared for Publication

in

The Journal of Physical Chemistry

Department of Chemistry
University of Rochester
Rochester, New York 14627

October 1984

Reproduction in whole or in part is permitted for any
purpose of the United States Government.

This document has been approved for public release
and sale; its distribution is unlimited.

84 11 21 052

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER UROCHESTER/DC/84/TR-56	2. GOVT ACCESSION NO. AD-A149 961	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Phase-Dissipative Mechanisms for Laser-Induced Surface Desorption/Dissociation Processes		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Xi-Yi Huang, Thomas F. George, Jian-Min Yuan and L. M. Narducci		8. CONTRACT OR GRANT NUMBER(s) N00014-80-C-0472
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Rochester Rochester, New York 14627		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 056-749
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Chemistry Program Code 472 Arlington, Virginia 22217		12. REPORT DATE October 1984
		13. NUMBER OF PAGES 11
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document has been approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Prepared for publication in the Journal of Physical Chemistry.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) ADMOLECULE-SURFACE SYSTEMS LASER-INDUCED VIBRATIONAL EXCITATION ENERGY-RELAXATION MECHANISMS PHASE-RELAXATION MECHANISMS T ₂ PROCESSES DESORPTION DISSOCIATION QUANTUM MECHANICAL STUDY ZWANZIG PROJECTORS WEIDLICH TECHNIQUE		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Vibrational excitation and relaxation for a molecule adsorbed on a surface is investigated in connection with desorption/dissociation processes. The population equations for a laser-driven anharmonic oscillator involving both energy- and phase-relaxation mechanisms are derived with the Zwanzig projector technique and eigenfunction-expansion method due to Weidlich. Phase relaxation is seen to assist in the excitation of the resonant active mode.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

PHASE-DISSIPATIVE MECHANISMS
FOR LASER-INDUCED SURFACE DESORPTION/DISSOCIATION PROCESSES

Xi-Yi Huang and Thomas F. George

Department of Chemistry

University of Rochester

Rochester, New York 14627

Jian-Min Yuan and L. M. Narducci

Department of Physics and Atmosphere Science

Drexel University

Philadelphia, Pennsylvania 19104



Accession For	
NTIS GRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Avail. and/or	
Dist	Special
A1	

Abstract

Vibrational excitation and relaxation for a molecule adsorbed on a surface is investigated in connection with desorption/dissociation processes. The population equations for a laser-driven anharmonic oscillator involving both energy- and phase-relaxation mechanisms are derived with the Zwanzig projector technique and eigenfunction-expansion method due to Weidlich. Phase relaxation is seen to assist in the excitation of the resonant active mode.

I. Introduction

Infrared laser radiation impinging onto a molecule adsorbed on a solid surface can resonantly excite the ad molecular internal vibrational/rotational modes, or deposit photon energy in the ad bond with which the molecule is bound to the surface.¹⁻⁵ It has been shown experimentally^{6,7} that SF_6 and CH_3F molecules adsorbed on NaCl surfaces at low temperatures can be desorbed by the resonant excitation of the adsorbate internal vibration, namely, the ν_3 mode. The treatment of the problem of many degrees of freedom for the molecule-surface system is very complicated and generally an insoluble problem. Interactions between the resonant active vibrational mode and the other inactive modes, and between the ad molecule and surface excitations - surface phonons, surface plasmons, reflected photons (i.e., photons emitted by the ad molecule and reflected by the surface), etc. - create a large number of dissipative channels by which absorption, dissociation, desorption or chemical reaction on the surface are strongly influenced. (Desorption is the breaking of the ad bond between the ad molecule and the surface, and dissociation is the breaking of a bond within the ad molecule.) In this Letter, we are especially interested in the influence of surface-dephasing relaxation on the excitation of the resonant mode, which plays an important role in the process of desorption/dissociation.

II. Theory

Under normal circumstances, due to the vast number of degrees of freedom of the ad molecule-surface system, the intrasystem relaxation can be very fast, and in fact, comparable to or faster than the rate of photon absorption. In this case, energy randomization or statistical thermodynamical behavior may at least partly occur in desorption/dissociation processes on surfaces.⁸ The IR-laser-driven

resonant active mode absorbs photons and transfers its energy to other vibrational modes or the surface, and then "interference" superposition of these many vibrations can instantaneously accumulate energy in the weakest bond to lead to molecule dissociation or desorption from the surface, if this energy is beyond the appropriate threshold.^{9,10} As an example, the threshold temperature for thermal dissociation of SF_6 in the gas phase is measured to be 1600 K.⁸ At this temperature, the average occupation number of the resonant mode (ν_3) of SF_6 is $\bar{n}=0.76$, which means that, while the energy deposited in the molecule has gone into the ν_3 mode to exceed the critical value for dissociation, the other modes remain in low excited states.

We shall confine ourselves here to the excitation and relaxation of the active anharmonic vibrational mode which is driven by a resonant coherent field and controlled by both energy (T_1) and phase (T_2) relaxation. For this purpose, we treat the many other non-active modes and surface excitations (i.e., phonons, plasmons, reflected photons, etc.) as a large thermal reservoir, by which dissipation mechanisms are provided. A reduced density operator W for the resonant mode obeys the master equation

$$\begin{aligned} \frac{dW}{dt} &= \left[\frac{dW}{dt} \right]_{\text{coherent}} + \left[\frac{dW}{dt} \right]_{\text{incoherent}} \\ &= \left[-iL_c + \Lambda_1 + \Lambda_2 \right] W. \end{aligned} \quad (1)$$

The reversible term is $-iL_c W \equiv -(i/\hbar)[H, W]$, where $H = H_M + H_F$ is the sum of anharmonic oscillator and external field interactions,¹¹

$$H_M = \hbar(\omega - \omega_L) a^\dagger a - \hbar \epsilon a^\dagger a (a^\dagger a + 1) \quad (2)$$

$$H_F = \mu E(t)(a^\dagger + a). \quad (3)$$

Here a^\dagger and a are oscillator ladder operators, μ is the molecular dipole moment, $E(t)$ is the applied electric field, ω_L is the laser frequency, and ϵ is a positive anharmonicity. The dissipative terms are given by^{12,13}

$$\begin{aligned} \Lambda_1 W = & \gamma_\downarrow([a, Wa^\dagger] + [aW, a^\dagger]) \\ & + \gamma_\uparrow([a^\dagger, Wa] + [a^\dagger W, a]), \end{aligned} \quad (4)$$

$$\Lambda_2 W = \eta([a^\dagger a, Wa^\dagger a] + [a^\dagger aW, a^\dagger a]). \quad (5)$$

$\Lambda_2 W$ is responsible for a pure dephasing in the decay of the off-diagonal matrix elements,¹⁴ and γ_\uparrow , γ_\downarrow and η contain implicit information about the reservoir, including the surface. With the dissipative terms [Eqs. (4) and (5) only], the relaxation behavior of the expectation values of the amplitude and number operator is governed by the equations $\frac{d}{dt}\langle a \rangle = -(\gamma_\downarrow - \gamma_\uparrow + \eta)\langle a \rangle$ and $\frac{d}{dt}\langle a^\dagger a \rangle = 2(\gamma_\downarrow - \gamma_\uparrow)\langle a^\dagger a \rangle + 2\gamma_\uparrow$. Λ_2 thus affects only the off-diagonal elements of the density operator by providing a pure phase-destroying term with a rate of η . In the Born approximation, the diagonal parts of the density matrix obey the equation

$$\frac{d}{dt} W_n = \langle n | \Lambda_1 W_r | n \rangle - \int_0^t d\tau \langle n | P L_i e^{-iL_M(t-\tau)} e^{\Lambda_2(t-\tau)} L_i W_r(\tau) | n \rangle, \quad (6)$$

where L_i is a Liouville operator due to the oscillator-field interaction, P is a projection operator used to obtain the diagonal part of the density operator,¹⁵ and $W_r \equiv PW$ is the relevant part of the density matrix.

The difficult problem here is the explicit evaluation of the matrix elements that appear inside the integral in Eq. (6). We use the Weidlich eigenfunction-expansion method^{12,13} to evaluate these elements. The "eigenstate" of the super-operators Λ_2 and Λ_2^\dagger are $||A_{pq}\rangle$ and $||B_{pq}\rangle$, respectively, and under the Markoff assumption, the "eigenvalues" of Λ_2 have the form $\lambda_{pq} \approx -\eta q^2$, where $p = 0, 1, 2, \dots$ and $q = 0, \pm 1, \pm 2, \dots$. The equations of motion for the diagonal parts of the density matrix are then given as

$$\dot{W}_n = \langle n | \dot{A}_1 W_r | n \rangle - \int_0^t d\tau \sum_{pq} e^{-nq^2(t-\tau)} \times \text{Tr}(B_{pq}^+ L_i W_r(\tau)) \langle n | L_i e^{-iL_M(t-\tau)} A_{pq} | n \rangle. \quad (7)$$

By a lengthy but straightforward algebraic calculation, we obtain a simpler set of coupled differential equations to describe the excitation and relaxation behavior for the resonant vibrational mode:

$$\dot{W}_n = 2\kappa [(n+1) W_{n+1} - n W_n] - 2\eta \left[\frac{\Omega_R(t)}{\eta} \right]^2 \left\{ \frac{n(W_n - W_{n-1})}{1 + [2\frac{\epsilon}{\eta}(n-1) - \frac{\Delta}{\eta}]^2} + \frac{(n+1)(W_n - W_{n+1})}{1 + (2\frac{\epsilon}{\eta}n - \frac{\Delta}{\eta})^2} \right\}. \quad (8)$$

The decay parameters κ ($\equiv \gamma_{\downarrow} - \gamma_{\uparrow}$) and η are determined by the internal mode-mode coupling and the ad molecule-surface interaction, and for a fixed temperature and ad molecule-surface distance they can be considered as constants (here a cold surface of 0 K has been assumed), $\Omega_R(t)$ is the Rabi frequency, and Δ is the detuning between the ad molecular fundamental and laser frequencies.

The first two terms with the factor of 2κ on the right-hand side of Eq. (8) describe the energy relaxation processes which always lead to population loss in the resonant mode. In the next terms with the factors of η describing the influence of the phase relaxation, we notice that there are competitive factors: upon an increase of the value of η , the phase relaxation tends to diminish the detuning and anharmonicity "bottleneck" effects, and the effective Rabi frequency $\Omega_R/\eta^{1/2}$ tends to decrease.

III. Calculations and Results

Figure 1 shows the influences of the phase relaxation. The time evolution for the average occupation number \bar{n} of a driven damped anharmonic oscillator is displayed, with consecutive values of the phase-relaxation parameter η ($\eta = 1, 5, 9, 13$). We note that there is a range of values of η which assist the excitation of the ad molecular active mode. For the optimal parameter $\eta = 5$ in our specific case, Figure 2 displays the time-dependent population distributions for several excited states, where considerable population is seen. One would generally expect the phase decay η to be much more enhanced than κ by the inactive adspecies vibrational modes and by the surface.¹⁶ The phase-dissipation mechanism may be generated in several ways in an ad molecule-surface system as follows: (a) intramolecular coupling between the active and inactive vibrational modes;¹⁷ (b) intermolecular collisions when the ad molecules can migrate on the surface;¹⁸ (c) rotational relaxation within the vibrational level manifold of the ad molecule;¹¹ (d) self-coupling of the ad molecule due to surface-reflected photons.^{19,20}

Acknowledgments

This research was supported in part by the Air Force Office of Scientific Research (AFSC), United States Air Force, under Grant AFOSR-82-0046, and the Office of Naval Research. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon. TFG acknowledges the Camille and Henry Dreyfus Foundation for a Teacher-Scholar Award (1975-84).

References

1. M. S. Slusky and T. F. George, Chem. Phys. Lett., 57, 474 (1978).
2. J. Lin and T. F. George, Surf. Sci., 100, 381 (1980).
3. J. Lin and T. F. George, J. Phys. Chem., 84, 2957 (1980).
4. D. Lucas and G. E. Ewing, Chem. Phys., 58, 385 (1981).
5. Z. W. Gortel, H. J. Kreuzer, P. Piercy and R. Teshima, Phys. Rev. B, 27, 5066 (1983).
6. J. Heidberg, H. Stein, A. Nestmann, E. Hoefs and I. Hussla, in Laser-Solid Interactions and Laser Processing, AIP Conf. Proc. 50, ed. by S. D. Ferris, H. J. Leamy and J. M. Poate (Am. Inst. Phys., New York, 1979), pp. 49-54.
7. J. Heidberg, H. Stein and E. Riehl, Phys. Rev. Lett. 49, 666 (1982); *ibid.*, Surf. Sci. 126, 183 (1983).
8. For a similar case in the gas phase, see J. G. Black, E. Yablonovitch, N. Bloembergen and S. Mukamel, Phys. Rev. Lett., 38, 1131 (1977).
9. N. B. Slater, Theory of Unimolecular Reactions (Cornell University, Ithaca, 1959).
10. P. J. Robinson and K. A. Holbrook, Unimolecular Reactions (Wiley, New York, 1972), Chapt. 2.
11. L. M. Narducci, S. S. Mitra, R. A. Shatas and C. A. Coulter, Phys. Rev. A, 16, 247 (1977).
12. W. Weidlich, Z. Phys., 241, 325 (1971).
13. X. Y. Huang, L. M. Narducci and J. M. Yuan, Phys. Rev. A 23, 3084 (1981).
14. A. H. Zewail, Acc. Chem. Res. 13, 360 (1980).
15. R. Zwanzig, Physica (Utrecht) 30, 1109 (1964).
16. P. L. Knight and P. W. Milonni, Physics Reports 66, 22 (1980).
17. J. Lin and T. F. George, Theor. Chim. Acta, 61, 243 (1983).
18. X. Y. Huang, J. D. Cresser and J. H. Eberly, J. Chem. Phys., in press.
19. X. Y. Huang, J. Lin and T. F. George, J. Chem. Phys., 80, 893 (1984).
20. X. Y. Huang and T. F. George, J. Phys. Chem., in press.

Figure Captions

1. Time dependence of the average occupation number $\bar{n} = \langle a^\dagger a \rangle$ for (cw) laser-driven anharmonic oscillator in contact with a bath. The curves correspond to $\eta = 1, 5, 9, 13$, and the others parameters are $(\kappa, \Omega_R, \epsilon, \Delta) = (0.01, 0.5, 1.5, 3.0)$, where all rates are in the unit of the transverse relaxation¹³ rate. This figure shows that when the vibrational linewidth η is smaller than the detuning Δ , the maximum vibrational excitation is small but grows as η increases. Excessively large values of η eventually reduce the average excitation level, presumably due to the spread of population within the linewidth.
2. The probability of occupation of level n , $W_n(t)$, as a function of time. η is fixed at 5, and the other parameters are given in Figure 1.

Figure 1





Figure 2



DL/413/83/01
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Naval Ocean Systems Center Attn: Technical Library San Diego, California 92152	1
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	Dean William Tolles Naval Postgraduate School Monterey, California 93940	1
Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. Vincent Schaper DTNSRDC Code 2830 Annapolis, Maryland 21402	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801	1
Dr. David L. Nelson Chemistry Program Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1		

TECHNICAL REPORT DISTRIBUTION LIST, 056

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Surface Chemistry Division (6170)
455 Overlook Avenue, S.W.
Washington, D.C. 20375

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. Chia-wei Woo
Department of Physics
Northwestern University
Evanston, Illinois 60201

Dr. Robert M. Hexter
Department of Chemistry
University of Minnesota
Minneapolis, Minnesota

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. Adolph B. Amster
Chemistry Division
Naval Weapons Center
China Lake, California 93555

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60201

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. S. L. Bernasek
Princeton University
Department of Chemistry
Princeton, New Jersey 08544

TECHNICAL REPORT DISTRIBUTION LIST, 056

Dr. F. Carter
Code 6132
Naval Research Laboratory
Washington, D.C. 20375

Dr. Richard Colton
Code 6112
Naval Research Laboratory
Washington, D.C. 20375

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Professor R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York ~~22217~~
12301

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 5570
Naval Research Laboratory
Washington, D.C. 20375

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
California Institute of Technology
Division of Chemistry and Chemical
Engineering
Pasadena, California 91125

Professor E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
Southampton University
Southampton SO9 5NH
Hampshire, England

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

Dr. Richard Smardzewski
Code 6130
Naval Research Laboratory
Washington, D.C. 20375

TECHNICAL REPORT DISTRIBUTION LIST, 056

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. P. Hansma
Physics Department
University of California
Santa Barbara, California 93106

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Professor T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
*BM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Professor Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Captain Lee Myers
AFOSR/NC
Bolling AFB
Washington, D.C. 20332

Professor Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Professor D. Hercules
University Pittsburgh
Chemistry Department
Pittsburgh, Pennsylvania 15260

Professor N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. G. D. Stein
Mechanical Engineering Department
Northwestern University
Evanston, Illinois 60201

Professor A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Professor G. H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. David Squire
Army Research Office
P.O. Box 12211
Research Triangle Park, NC 27709

